


WHAT IS CLAIMED IS:

1. A compound of Formula I:



Formula I

or a pharmaceutically acceptable salt thereof, wherein:

x is 1, 2 or 3;

R represents from 0 to 4 substituents independently chosen from halogen, hydroxy, optionally substituted alkoxy, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, cyano, amino, nitro, -COOH, carboxamide, optionally substituted mono- and di-alkyl amino, optionally substituted haloalkyl, and optionally substituted haloalkoxy;

R¹ is selected from alkyl, alkenyl, alkynyl, cycloalkyl, (cycloalkyl)alkyl, aryl, heteroaryl, (aryl)alkyl, (heteroaryl)alkyl, and indanyl, each of which is optionally substituted;

R², R³ and each occurrence of R⁴ are independently selected from hydrogen, halogen, optionally substituted alkyl, and optionally substituted alkoxy;

R⁵ and R⁶ are independently selected from

- (i) hydrogen, halogen, hydroxy, amino, and cyano; and
- (ii) alkyl, alkenyl, alkynyl, alkoxy, haloalkyl, haloalkoxy, and mono- and di-(alkyl)amino, each of which is optionally substituted;


R⁷ is: (a) (i) hydrogen; or (ii) alkyl, alkenyl, alkynyl, alkoxy or arylalkyl, each of which is optionally substituted; and

Ar¹ is:

- (i) phenyl, naphthyl, biphenyl, or heterocycle, each of which is optionally substituted; or
- (ii) optionally substituted phenyl fused to a 5- to 7-membered saturated or partially unsaturated ring having from 5 to 7 ring atoms, with 0, 1, or 2 ring atoms independently chosen from N, O and S, and with remaining ring atoms being carbon;

or

(b) taken together with Ar¹ and the carbon atom to which R⁷ and Ar¹ are attached to form an optionally substituted group of the formula:



wherein p is an integer from 1 to about 3; and

Ar² is (i) optionally substituted aryl or (ii) optionally substituted heteroaryl having 5 to 7 ring atoms and from 1 to 3 ring heteroatoms independently selected from N, O and S.

2. A compound or salt according to Claim 1, wherein:

x is 1;

R represents from 0 to 4 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, -COOH, carboxamide, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, mono- and di-(C₁-C₆alkyl)amino, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy;

R¹ is selected from (aryl)C₀-C₆alkyl, (heteroaryl)C₀-C₆alkyl, and indanyl, each of which is substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, carboxamide, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, mono- and di-(C₁-C₆alkyl)amino, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy;

R², R³, and each occurrence of R⁴ are independently selected from hydrogen, halogen, C₁-C₆alkyl, and C₁-C₆alkoxy;

R⁵ and R⁶ are independently selected from hydrogen, halogen, cyano, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆alkoxy, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, hydroxy, amino, and mono- and di-(C₁-C₆alkyl)amino;

and either:


(a) R⁷ is (i) hydrogen; or (ii) C₁-C₆alkyl, C₂-C₆alkenyl, C₁-C₆alkynyl, C₁-C₆alkoxy or (aryl)C₁-C₆alkyl, each of which is optionally substituted; and

Ar¹ is (i) phenyl; (ii) naphthyl; (iii) biphenyl; (iv) a heterocyclic group having 1 or 2 rings, 3 to 8 atoms in each ring and in at least one ring from 1 to 3 heteroatoms independently selected from N, O and S; or (v) phenyl fused to a 5- to 7-membered saturated or partially unsaturated ring having from 5 to 7 ring atoms, with 0, 1 or 2 ring atoms chosen from N, O and S, and with remaining ring atoms being carbon;

wherein each of (i), (ii), (iii), (iv) and (v) is substituted with from 0 to 4 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, C₁-C₆alkoxy, C₁-

C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆alkoxycarbonyl, -COOH, carboxamide, mono- and di-(C₁-C₆alkyl)amino, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy; or

(b) R⁷ is taken together with Ar¹ and the carbon atom to which R⁷ and Ar¹ are attached to form a group of the formula:



, substituted with from 0 to 4 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, -COOH, carboxamide, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, mono- and di-(C₁-C₆alkyl)amino, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy, wherein p is an integer from 1 to about 3; and

Ar² is aryl or heteroaryl, each of which is substituted with from 0 to 5 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆alkoxycarbonyl, carboxamide, mono- and di-(C₁-C₆alkyl)carboxamide, mono- and di-(C₁-C₆alkyl)amino, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy.

3. A compound or salt according to Claim 1, wherein R¹ is indanyl, substituted with 0, 1, or 2 substituents independently selected from halogen, hydroxy, C₁-C₂alkoxy, C₁-C₂alkyl, haloC₁-C₂alkyl, and haloC₁-C₂alkoxy.

4. A compound or salt according to Claim 1, wherein R¹ is phenyl(C₀-C₄alkyl), pyridyl(C₀-C₄alkyl), C₀-C₄alkyl, or indolyl(C₀-C₄alkyl), each of which is substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, C₁-C₂alkoxy, C₁-C₂alkyl, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy.

5. A compound or salt according to Claim 1, wherein R¹ is phenyl(C₀-C₂alkyl) substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, C₁-C₂alkoxy, C₁-C₂alkyl, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy.

6. A compound or salt according to Claim 1, wherein R² and R³ are hydrogen.


7. A compound or salt according to Claim 1, wherein each R^4 is independently hydrogen or C_1 - C_6 alkyl.

8. A compound or salt according to Claim 1, wherein R^5 and R^6 are independently selected from hydrogen, halogen, C_1 - C_2 alkyl and C_1 - C_2 alkoxy.

9. A compound or salt according to Claim 1, wherein R represents 0, 1, or 2 substituents independently selected from C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_2 haloalkyl, C_1 - C_2 haloalkoxy, fluoro, and chloro.

10. A compound or salt according to Claim 1, wherein
R represents 0, 1, or 2 substituents independently selected from hydrogen, methyl, ethyl, methoxy, trifluoromethyl, trifluoromethoxy, fluoro, and chloro;
 R^2 , R^3 , and R^6 are hydrogen; and
 R^5 , R^7 , and each R^4 are independently selected from hydrogen, methyl, and ethyl.

11. A compound or salt according to Claim 1, of Formula II



Formula II

wherein:

R represents from 0 to 4 substituents independently chosen from fluoro, chloro, hydroxy, optionally substituted C_1 - C_6 alkoxy and optionally substituted C_1 - C_6 alkyl;
 R^4 is hydrogen, optionally substituted C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, fluoro, or chloro;
 R^5 and R^6 are independently selected from hydrogen, fluoro, chloro, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, C_1 - C_6 haloalkyl, and C_1 - C_6 haloalkoxy; and
 R^7 is hydrogen or C_1 - C_6 alkyl.

12. A compound or salt according to Claim 11, wherein

R represents from 0 to 4 substituents independently chosen from fluoro, chloro, hydroxy, C₁-C₆alkoxy, and C₁-C₆alkyl;

R¹ is selected from C₃-C₇cycloalkyl, (C₃-C₇cycloalkyl)C₁-C₄alkyl, (heteroaryl)C₀-C₄alkyl, (aryl)C₀-C₄alkyl, and indanyl, each of which is substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, -COOH, carboxamide, mono- and di-(C₁-C₆alkyl)amino, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy;

R⁴ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, fluoro, or chloro;

R⁵ and R⁶ are independently selected from hydrogen, fluoro, chloro, C₁-C₆alkyl, C₁-C₆alkoxy, haloC₁-C₆alkyl, and C₁-C₆haloalkoxy; and

R⁷ is hydrogen or C₁-C₆ alkyl.

13. A compound or salt according to Claim 11 or 12, wherein

Ar¹ is:

(i) phenyl substituted with from 0 to 4 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, carboxamide, C₁-C₃alkoxy, C₁-C₃alkyl, C₁-C₂haloalkyl, C₁-C₂alkoxycarbonyl, mono- and di-(C₁-C₂alkyl)amino, and C₁-C₂haloalkoxy;

(ii) naphthyl;

(iii) heterocyclic groups having 1 or 2 rings, 3 to 8 atoms in each ring and in at least one ring from 1 to 3 heteroatoms independently selected from N, O and S;

(iv) biphenyl, wherein each phenyl group is substituted with 0 to 2 groups independently selected from halogen, C₁-C₂alkyl, and C₁-C₂alkoxy; or

(v) phenyl fused to a 5- to 7-membered saturated or partially unsaturated ring having from 5 to 7 ring atoms, with 0, 1, or 2 ring atoms independently chosen from N, O and S, and with remaining ring atoms being carbon;

wherein each of (ii), (iii), (iv) and (v) is substituted with from 0 to 4 substituents independently selected from halogen, hydroxy, C₁-C₂alkyl, C₁-C₂alkoxy, C₁-C₂haloalkyl, and haloC₁-C₂alkoxy.

14. A compound or salt according to Claim 13, wherein Ar² is phenyl or heteroaryl having about 5 to 7 ring atoms and between 1 and 3 ring heteroatoms independently selected from N, O and S, each of which is substituted with from 0 to 5 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, C₁-C₃alkoxy, C₁-C₃alkyl, carboxamide, dimethylcarboxamide, mono- and di-(C₁-C₂alkyl)amino, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy.

15. A compound or salt according to Claim 11, wherein Ar² is phenyl or heteroaryl having about 5 to 7 ring atoms and between 1 and 3 ring heteroatoms independently selected from N, O and S, each of which is substituted with from 0 to 5 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, C₁-C₃alkoxy, C₁-C₃alkyl, carboxamide, dimethylcarboxamide, mono- and di-(C₁-C₂alkyl)amino, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy.

16. A compound or salt according to claim 12, wherein:

R represents from 0 to 2 substituents independently chosen from fluoro, chloro, hydroxy, methoxy, ethoxy, methyl, and ethyl;

R¹ is 1-indanyl or 2-indanyl, each of which is substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, -COOH, carboxamide, mono- and di-(C₁-C₆alkyl)amino, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy;

R⁷ is hydrogen, methyl or ethyl;

Ar₁ is

- (i) phenyl substituted with from 0 to 5 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, carboxamide, C₁-C₃alkoxy, C₁-C₃alkyl, C₁-C₂alkoxycarbonyl, mono- and di-(C₁-C₂alkyl)amino, haloC₁-C₂alkyl, and haloC₁-C₂alkoxy;
- (ii) naphthyl;
- (iii) heterocyclic groups having 1 or 2 rings, 3 to 8 atoms in each ring, and in at least one ring from 1 to 3 heteroatoms independently selected from N, O and S;
- (iv) biphenyl; or

(v) phenyl fused to a 5- to 7-membered saturated or partially unsaturated ring having from 5 to 7 ring atoms, with 0, 1, or 2 ring atoms independently chosen from N, O and S, and with remaining ring atoms being carbon;

wherein each of (ii), (iii), (iv) and (v) is substituted with from 0 to about 4 substituents independently selected from halogen, hydroxy, C₁-C₂alkyl, C₁-C₂alkoxy, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy; and

Ar² is phenyl, pyridyl, thiazolyl, pyrimidyl, pyridizynyl, imidazolyl, oxazolyl, isoxazolyl and triazolyl, each of which is substituted with from 0 to 3 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, -COOH, C₁-C₃alkoxy, C₁-C₃alkyl, carboxamide, dimethylcarboxamide, mono- and di-(C₁-C₂alkyl)amino, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy.

17. A compound or salt according to claim 16, wherein

R represents from 0 to 2 substituents independently chosen from fluoro, chloro, hydroxy, methoxy, ethoxy, methyl and ethyl;

R¹ is 2-indanyl, substituted with 0, 1, or 2 substituents independently selected from fluoro, chloro, hydroxy, methyl, ethyl, methoxy, ethoxy, mono-, di- and tri-fluoromethyl, and mono-, di-, and tri-fluoromethoxy;

R⁴ is hydrogen, fluoro, chloro, methyl, ethyl, methoxy, mono-, di-, or tri-fluoromethyl, or mono-, di- or tri-fluoromethoxy;

R⁵ and R⁶ are independently selected from hydrogen, fluoro, chloro, methyl, ethyl, methoxy, ethoxy, mono-, di- and tri-fluoromethyl, and mono-, di-, and tri-fluoromethoxy;

Ar¹ is:

(i) phenyl, substituted with from 0 to 3 substituents independently selected from fluoro, chloro, bromo, hydroxy, methyl, methoxy, ethyl, ethoxy, mono-, di- and tri-fluoromethyl, and mono-, di-, and tri-fluoromethoxy; or

(ii) naphthyl, substituted with from 0 to 3 substituents independently selected from fluoro, chloro, hydroxy, methyl, ethyl, methoxy, and ethoxy; and

Ar² is phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-thienyl, 3-thienyl or 1,3-thiazol-2-yl, each of which is substituted with from 0 to 3 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, -COOH, C₁-C₃alkoxy, C₁-C₃alkyl, carboxamide,

dimethylcarboxamide, mono- and di-(C₁-C₂alkyl)amino, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy.

18. A compound or salt according to Claim 12, wherein:

R represents from 0 to 2 substituents independently chosen from fluoro, chloro, hydroxy, methoxy, ethoxy, methyl, and ethyl;

R¹ is phenyl(C₀-C₂alkyl), substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cyano, amino, nitro, -COOH, carboxamide, mono- and di-(C₁-C₆alkyl) amino, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy;

R⁷ is hydrogen, methyl, or ethyl;

Ar₁ is:


- (i) phenyl substituted with from 0 to 5 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, carboxamide, C₁-C₃alkoxy, C₁-C₃alkyl, C₁-C₂alkoxycarbonyl, mono- and di-(C₁-C₂alkyl)amino, C₁-C₂haloalkyl, and C₁-C₂haloC₁-C₂alkoxy;
- (ii) naphthyl;
- (iii) a heterocyclic group having 1 or 2 rings, 3 to 8 atoms in each ring, and in at least one ring from 1 to 3 heteroatoms independently selected from N, O and S;
- (iv) biphenyl; or
- (v) phenyl fused to a 5- to 7-membered saturated or partially unsaturated ring having from 5 to 7 ring atoms, with 0, 1 or 2 ring atoms independently chosen from N, O and S, and with remaining ring atoms being carbon;

wherein each of (ii), (iii), (iv) and (v) is substituted with from 0 to about 4 substituents independently selected from halogen, hydroxy, C₁-C₂alkyl, C₁-C₂alkoxy, haloC₁-C₆alkyl, and haloC₁-C₂alkoxy; and

Ar² is phenyl, pyridyl, thiazolyl, pyrimidyl, pyridizynyl, imidazolyl, oxazolyl, isoxazolyl or triazolyl, each of which is substituted with from 0 to 3 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, -COOH, C₁-C₃alkoxy, C₁-C₃alkyl, carboxamide, dimethylcarboxamide, mono- and di-(C₁-C₂alkyl)amino, haloC₁-C₂alkyl and haloC₁-C₂alkoxy.

19. A compound or salt according to Claim 18, wherein R^1 is phenyl(C_0 - C_1 alkyl), substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 alkyl, $-COOH$, carboxamide, mono- and di-(C_1 - C_4 alkyl) amino, C_1 - C_2 haloalkyl, and C_1 - C_2 haloalkoxy.

20. A compound or salt according to Claim 1, of Formula III



Formula III

wherein:

R represents from 0 to 4 substituents independently chosen from fluoro, chloro, hydroxy, C_1 - C_6 alkoxy, and C_1 - C_6 alkyl;

R^4 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, fluoro, or chloro;

R^8 represents from 0 to 4 substituents independently chosen from C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, fluoro, and chloro;


R^5 and R^6 are independently selected from hydrogen, fluoro, chloro, C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, and halo C_1 - C_6 alkyl;

R^7 is hydrogen or C_1 - C_6 alkyl;

R^9 represents from 0 to 5 substituents independently chosen from halogen, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 haloalkyl, and C_1 - C_6 haloalkoxy; and

Ar^2 is (i) phenyl or (ii) heteroaryl having 5 to 7 ring atoms and from 1 to 3 ring heteroatoms independently selected from N, O and S, wherein each of (i) and (ii) is optionally substituted with from 1 to 5 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, $-COOH$, carboxamide, dimethylcarboxamide, C_1 - C_3 alkoxy, C_1 - C_3 alkyl, mono- and di-(C_1 - C_2 alkyl)amino, C_1 - C_2 haloalkyl, and C_1 - C_2 haloalkoxy.

21. A compound or salt according to Claim 1, of Formula IV



Formula IV

wherein:

R represents from 0 to 4 substituents independently chosen from fluoro, chloro, hydroxy, C₁-C₆alkoxy, and C₁-C₆alkyl;

R⁴ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, fluoro, or chloro;

R⁸ represents from 0 to 4 substituents independently chosen from C₁-C₆alkyl, C₁-C₆haloalkyl, fluoro, and chloro;

R⁵ and R⁶ are independently selected from hydrogen, fluoro, chloro, C₁-C₆alkyl, optionally substituted C₁-C₆alkoxy, and C₁-C₆haloalkyl;

R⁷ is hydrogen or C₁-C₆alkyl;


R¹⁰ represents from 0 to 5 substituents independently chosen from fluoro, chloro, bromo, iodo, hydroxy, nitro, cyano, -COOH, carboxamide, dimethylcarboxamide, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy; and

Ar₁ is:

- (i) phenyl substituted with from 0 to 5 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, carboxamide, C₁-C₃alkoxy, C₁-C₃alkyl, C₁-C₂haloalkyl, C₁-C₂alkoxycarbonyl, mono- and di-(C₁-C₂alkyl)amino, and C₁-C₂haloalkoxy;
- (ii) naphthyl;
- (iii) heterocyclic groups having 1 or 2 rings, 3 to 8 atoms in each ring and in at least one ring from 1 to 3 heteroatoms independently selected from N, O and S;
- (iv) biphenyl; or

- (v) phenyl fused to a 5- to 7-membered saturated or partially unsaturated ring having from 5 to 7 ring atoms, with 0, 1, or 2 ring atoms independently chosen from N, O and S, and with remaining ring atoms being carbon;
 wherein each of (ii), (iii), (iv) and (v) is substituted with from 0 to 4 substituents independently selected from halogen, hydroxy, C₁-C₂alkyl, C₁-C₂alkoxy, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy.

22. A compound or salt according to Claim 21, of Formula V



Formula V

wherein:

R represents from 0 to 4 substituents independently chosen from fluoro, chloro, hydroxy, methyl, ethyl, methoxy, or ethoxy;


R⁴ is hydrogen, fluoro, chloro, methyl, ethyl, methoxy, mono-, di- or tri-fluoromethyl, or mono-, di-, or tri-fluoromethoxy;

R⁵ and R⁶ are independently chosen from hydrogen, fluoro, chloro, methyl, methoxy, mono-, di- and tri-fluoromethyl, and mono-, di- and tri-fluoromethoxy;

R⁷ is hydrogen, methyl, or ethyl; and

R¹⁰ is hydrogen, fluoro, chloro, bromo, hydroxy, methyl, ethyl, methoxy, or ethoxy.

23. A compound of the Formula VI



Formula VI

or a pharmaceutically acceptable salt thereof, wherein:

R is hydrogen, fluoro, chloro, hydroxy, methyl, or methoxy;

R⁴ is hydrogen, methyl, or ethyl;

R⁸ is hydrogen, fluoro, chloro, methyl, or methoxy;


R⁹ is fluoro, chloro, methyl, ethyl, methoxy, ethoxy, mono-, di- or tri-fluoromethyl, or mono-, di-, or tri-fluoromethoxy;

R^{9A} represents 0, 1, or 2 substituents independently selected from hydrogen, fluoro, chloro, methyl, methoxy, mono-, di- and tri-fluoromethyl, and mono-, di-, and tri-fluoromethoxy;

R¹⁰ is hydrogen, fluoro, chloro, hydroxy, methyl, methoxy, mono-, di-, or tri-fluoromethyl, or mono-, di-, or tri-fluoromethoxy; and


R^{10A} represents from 0 to 3 substituents independently selected from hydrogen, fluoro, chloro, hydroxy, methyl, methoxy, mono-, di- and tri-fluoromethyl, and mono-, di-, and tri-fluoromethoxy.

24. A compound or salt according to claim 23, of Formula VII



Formula VII

25. A compound of Formula VIII



Formula VIII

or a pharmaceutically acceptable salt thereof, wherein:

A is N or CR¹⁰;

B is N or CR^{10A}, wherein at least one of A and B is not N;

R is hydrogen, fluoro, chloro, hydroxy, methyl, or methoxy;

R⁴ is hydrogen, methyl, or ethyl;


R⁸ is hydrogen, fluoro, chloro, methyl or methoxy;

R¹⁰, if present, is hydrogen, fluoro, chloro, hydroxy, nitro, cyano, methyl, methoxy, mono-, di- or tri-fluoromethyl, or mono-, di-, or tri-fluoromethoxy;

R^{10A}, if present, is hydrogen, fluoro, chloro, hydroxy, nitro, cyano, methyl, methoxy, mono-, di- or tri-fluoromethyl, or mono-, di-, or tri-fluoromethoxy; and

R^{10B} represents from 0 to 3 substituents independently selected from fluoro, chloro, hydroxy, nitro, cyano, methyl, methoxy, mono-, di- and tri-fluoromethyl, and mono-, di-, and tri-fluoromethoxy.

26. A compound or salt according to Claim 25, of Formula IX



Formula IX

wherein:


R is hydrogen, fluoro, chloro, hydroxy, methyl or methoxy;

R^4 is hydrogen, methyl, or ethyl;

R^8 is hydrogen, fluoro, chloro, methyl or methoxy; and

R^{10} is hydrogen, fluoro, chloro, hydroxy, methyl, methoxy, mono-, di- or tri-fluoromethyl, or mono-, di-, or tri-fluoromethoxy.

27. A compound according to Claim 1, of Formula X



Formula X

or a pharmaceutically acceptable salt thereof, wherein:

R is hydrogen, fluoro, chloro, hydroxy, methyl or methoxy;


R⁴ is hydrogen, methyl, or ethyl;

R⁸ is hydrogen, fluoro, chloro, methyl or methoxy;

R¹⁰ is hydrogen, fluoro, chloro, hydroxy, methyl, methoxy, mono-, di- or tri-fluoro methyl, or mono-, di- or tri-fluoromethoxy; and

R^{10A} represents from 0 to 3 substituents independently selected from hydrogen, fluoro, chloro, hydroxy, methyl, methoxy, mono-, di- and tri-fluoromethyl, and mono-, di-, and tri-fluoromethoxy.

28. A compound or salt according to Claim 1, of Formula XI



Formula XI

wherein:

R represents from 0 to 4 substituents independently chosen from fluoro, chloro, hydroxy, C₁-C₆alkoxy, and C₁-C₆alkyl;

R⁴ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, fluoro, or chloro;

R⁸ represents from 0 to 4 substituents independently chosen from C₁-C₆alkyl, C₁-C₆haloalkyl, fluoro, and chloro;

R⁵ and R⁶ are independently selected from hydrogen, fluoro, chloro, C₁-C₆alkyl, optionally substituted C₁-C₆alkoxy, and C₁-C₆haloalkyl;

R⁷ is hydrogen or C₁-C₆alkyl;

R¹⁰ represents from 0 to 5 substituents independently chosen from fluoro, chloro, bromo, iodo, hydroxy, nitro, cyano, -COOH, carboxamide, dimethylcarboxamide, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy; and

Ar₁ is:

- (i) phenyl optionally substituted with from 1 to 5 substituents independently selected from halogen, hydroxy, cyano, amino, nitro, -COOH, carboxamide, C₁-C₃alkoxy, C₁-C₃alkyl, C₁-C₂haloalkyl, C₁-C₂alkoxycarbonyl, mono- and di-(C₁-C₂alkyl)amino, and C₁-C₂haloalkoxy;
 - (ii) naphthyl;
 - (iii) heterocyclic groups having 1 or 2 rings, 3 to 8 atoms in each ring and in at least one ring from 1 to 3 heteroatoms independently selected from N, O and S;
 - (iv) biphenyl; or
 - (v) phenyl fused to a 5- to 7-membered saturated or partially unsaturated ring having from 5 to 7 ring atoms, with 0, 1, or 2 ring atoms independently chosen from N, O and S, and with remaining ring atoms being carbon;
- wherein each of (ii), (iii), (iv) and (v) is optionally substituted with from 1 to 4 substituents independently selected from halogen, hydroxy, C₁-C₂alkyl, C₁-C₂alkoxy, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy.

29. A compound according to Claim 1, which is:

N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[8-methoxy-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;

N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-methyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;

N-(indan-2-yl)-2-[1-ethyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)acetamide;

2-[1-(2,4-difluorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;

N-(2-fluorobenzyl)-N-{3-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-3-oxopropyl}indan-2-amine;

N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanamide;

N-(indan-2-yl)-2-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(pyridin-2-ylmethyl)acetamide;

N-(indan-2-yl)-2-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(pyridin-3-ylmethyl)acetamide;

2-[1-(2-bromophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;

N-(indan-2-yl)-2-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(1,3-thiazol-2-ylmethyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-(2-methoxyphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-2-[1-(2,3-dimethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R)-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(4R)-4-methyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1S)-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-2-[1-(2-ethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)acetamide;
 N-(2-Fluoro-benzyl)-N-indan-2-yl-2-(4-methyl-1-o-tolyl-3,4-dihydro-1H-isoquinolin-2-yl)-acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R,4S)-4-methyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1S,4S)-4-methyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 2-{[1-(indan-2-yl)-2-phenyl-1H-imidazol-5-yl]methyl}-1-(2-methylphenyl)-1,2,3,4-tetrahydroisoquinoline;
 (2S)-N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R)-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanamide;
 N-(indan-2-yl)-2-[1-(3,4-dimethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)acetamide;
 2-[1-(2,3-dichlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-[4-fluoro-2-(trifluoromethyl)phenyl]-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-[5-fluoro-2-(trifluoromethyl)phenyl]-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 (2S)-N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1S)-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanamide;
 2-[(1S)-1-(2-bromophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;

N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1S)-1-[2-(trifluoromethyl)phenyl]-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 (2S)-N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R)-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanamide;
 2-[1-(1,1'-biphenyl-2-yl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-(1-naphthyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R,4R)-4-methyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 2-[1-(2-chloro-3-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-(3-fluoro-2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-2-[1-(2,5-dimethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-[3-(trifluoromethyl)phenyl]-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 2-[1-(5-chloro-2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 2-[1-(2-chloro-5-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 2-[1-(2,3-dihydro-1-benzofuran-7-yl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R,4R)-1-(2-fluorophenyl)-4-methyl-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(3S)-3-methyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-2-[1-(2,6-dimethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-(5-fluoro-2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R)-1-(2-fluorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 2-[(1R,4R)-1-(2-chlorophenyl)-4-methyl-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 (2S)-N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R)-1-(2-fluorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanamide;

N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-(3-fluoro-4-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-(1-quinolin-8-yl-3,4-dihydroisoquinolin-2(1H)-yl)acetamide;
 (2S)-N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R)-1-(1-naphthyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanamide;
 (2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)propanamide;
 2-[1-(3-chloro-2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-(2-fluoro-5-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 (2S)-N-(indan-2-yl)-2-[(1R)-1-(2,3-dimethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)propanamide;
 (2S)-N-(indan-2-yl)-2-[(1S)-1-(2,3-dimethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)propanamide;
 (2S)-N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R)-1-[2-(trifluoromethyl)phenyl]-3,4-dihydroisoquinolin-2(1H)-yl]propanamide;
 (2S)-2-[(1R)-1-(2-bromophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)propanamide;
 (2S)-N-(indan-2-yl)-2-[(1R)-1-(2,6-dimethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)propanamide;
 N-(indan-2-yl)-2-[6,7-dimethyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)acetamide;
 N-(indan-2-yl)-2-[7,8-dimethyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)acetamide;
 2-[1-(2,3-difluorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 (2S)-2-[(1R)-1-(2,3-dihydro-1-benzofuran-7-yl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)propanamide;
 methyl 4-(2-{2-[indan-2-yl(2-fluorobenzyl)amino]-2-oxoethyl}-1,2,3,4-tetrahydroisoquinolin-1-yl)benzoate;
 (2S)-N-benzyl-2-[(1R)-1-(2-bromophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-chloro-4-hydroxybenzyl)propanamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[8-methyl-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 N-(indan-2-yl)-N-(3-methoxybenzyl)-2-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;

N-(indan-2-yl)-N-(2-fluorobenzyl)-2-(1-mesityl-3,4-dihydroisoquinolin-2(1H)-yl)acetamide;
 2-[1-(2,6-difluorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 N-(2-fluorobenzyl)-2-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-phenylacetamide;
 N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-(2-methyl-1-naphthyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 (2S)-2-[(1R)-1-(2-chloro-5-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)propanamide;
 (2S)-N-(indan-2-yl)-2-[(1S)-1-(2,5-dimethylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)propanamide;
 (2S)-N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[(1R)-1-(2-fluoro-5-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanamide;
 (2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N,N-bis(2-fluorobenzyl)propanamide;
 4-[(indan-2-yl){[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetyl}amino)methyl]benzoic acid;
 (2S)-2-[(1R)-1-(2,6-difluorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)propanamide;
 (2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-hydroxybenzyl)propanamide;
 (2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)-N-(2-phenylethyl)propanamide;
 N-(indan-2-yl)-N-(3-hydroxybenzyl)-2-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 2-[1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)-N-(5-methoxy-indan-2-yl)acetamide;
 N-(2-fluorobenzyl)-N-(5-methoxy-indan-2-yl)-2-[1-[2-(trifluoromethyl)phenyl]-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;
 2-[1-(2,6-dichlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 2-[1-(2-chloro-6-fluorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;
 (2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(1H-imidazol-4-ylmethyl)propanamide;
 3-[[{(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanoyl}(indan-2-yl)amino)methyl]benzoic acid;

N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[1-[2-fluoro-6-(trifluoromethyl)phenyl]-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;

N-(indan-2-yl)-N-(2-fluorobenzyl)-2-[8-fluoro-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;

N-(indan-2-yl)-N-(2-fluoro-5-hydroxybenzyl)-2-[1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;

(2S)-2-[(1R)-1-(2,6-dichlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)propanamide;

(2S)-2-[(1R)-1-(2-chloro-6-fluorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)propanamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-[2-(2-fluorophenyl)ethyl]-N-(4-hydroxybenzyl)propanamide;

3-[[[(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]propanoyl](indan-2-yl)amino]methyl]-N,N-dimethylbenzamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(4-hydroxy-3,5-dimethylbenzyl)propanamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)propanamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-[2-(2-fluorophenyl)ethyl]-N-(2-phenylethyl)propanamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(2-fluorobenzyl)-N-[2-(2-fluorophenyl)ethyl]propanamide;

(2S)-2-[(1R)-1-(2,6-dichlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(4-hydroxy-3,5-dimethylbenzyl)propanamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(3-cyanobenzyl)-N-(indan-2-yl)propanamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(3-nitrobenzyl)propanamide;


N-(indan-2-yl)-N-(3-hydroxybenzyl)-2-[(1S)-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]acetamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(3-hydroxybenzyl)propanamide;

(2S)-2-[(1R)-1-(2-chlorophenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluoro-3-hydroxybenzyl)propanamide;

2-[8-chloro-1-(2-methylphenyl)-3,4-dihydroisoquinolin-2(1H)-yl]-N-(indan-2-yl)-N-(2-fluorobenzyl)acetamide;

31. A compound of Formula XII



Formula XII

or a pharmaceutically acceptable salt thereof, wherein:

R represents from 0 to 4 substituents independently chosen from fluoro, chloro, hydroxy, C₁-C₆alkoxy, C₁-C₆alkyl, C₁-C₂haloalkyl, and C₁-C₂haloalkoxy;

R¹ and Ar² are independently chosen from:

(i) phenyl(C₀-C₁alkyl), substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cyano, amino, nitro, -COOH, carboxamide, mono- and di-(C₁-C₆alkyl)amino, C₁-C₆haloalkyl and C₁-C₆haloalkoxy; and

(ii) 2-indanyl, substituted with 0, 1 or 2 substituents independently selected from fluoro, chloro, hydroxy, methyl, ethyl, methoxy, ethoxy, mono-, di- and tri-fluoromethyl, and mono-, di- and tri-fluoromethoxy;

R⁴ is C₁-C₆alkyl, C₁-C₂haloalkyl, fluoro or chloro;

R⁷ is hydrogen or C₁-C₆alkyl;

R⁹ represents from 0 to 5 substituents independently chosen from hydrogen, halogen, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy; and

|| represents a single or double bond.

32. A compound according to claim 31, wherein the compound is:

N-(2-Fluoro-benzyl)-N-indan-2-yl-2-(3-methyl-2-o-tolyl-piperidin-1-yl)-acetamide;

N-(2-Fluoro-benzyl)-N-indan-2-yl-2-(2-o-tolyl-piperidin-1-yl)-acetamide;


N-(2-Fluoro-benzyl)-N-indan-2-yl-2-[2-(2-methoxy-phenyl)-piperidin-1-yl]-acetamide;

N-(2-Fluoro-benzyl)-N-indan-2-yl-2-(2-o-tolyl-piperidin-1-yl)-propionamide.

2-(4,5-Dimethyl-6-phenyl-3,6-dihydro-2H-pyridin-1-yl)-N-(2-fluoro-benzyl)-N-indan-2-yl-acetamide;

2-(4,5-Dimethyl-6-o-tolyl-3,6-dihydro-2H-pyridin-1-yl)-N-(2-fluoro-benzyl)-N-indan-2-yl-acetamide;
or a pharmaceutically acceptable salt thereof.

33. A compound of Formula XIII



Formula XIII

or a pharmaceutically acceptable salt thereof, wherein:

R^1 and Ar^2 are independently chosen from:

- (i) phenyl(C_0 - C_1 alkyl), substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, C_1 - C_6 alkoxy, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cyano, amino, nitro, -COOH, carboxamide, mono- and di-(C_1 - C_6 alkyl) amino, C_1 - C_6 haloalkyl, and C_1 - C_6 haloalkoxy; and
- (ii) 2-indanyl, substituted with 0, 1 or 2 substituents independently selected from fluoro, chloro, hydroxy, methyl, ethyl, methoxy, ethoxy, mono-, di- and tri-fluoromethyl, and mono-, di- and tri-fluoromethoxy;


R^4 is C_1 - C_6 alkyl, halo C_1 - C_6 alkyl, fluoro or chloro;

R^3 and R^7 are independently hydrogen or C_1 - C_6 alkyl; and

R^9 and R^{10} independently represent from 0 to 5 substituents independently chosen from hydrogen, halogen, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 haloalkyl, and C_1 - C_6 haloalkoxy.

34. A compound according to claim 33, wherein the compound is (Benzhydryl-methyl-amino)-N-(2-fluoro-benzyl)-N-indan-2-yl-acetamide.

35. A compound of Formula XIV



Formula XIV

or a pharmaceutically acceptable salt thereof, wherein:

R represents from 0 to 2 substituents independently chosen from fluoro, chloro, hydroxy, C₁-C₆alkoxy, C₁-C₆alkyl, C₁-C₂alkyl, and C₁-C₂alkoxy;

R¹ and Ar² are independently chosen from:

(i) phenyl(C₀-C₁alkyl), substituted with from 0 to 3 substituents independently selected from halogen, hydroxy, C₁-C₆alkoxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cyano, amino, nitro, -COOH, carboxamide, mono- and di-(C₁-C₆alkyl) amino, haloC₁-C₆alkyl and haloC₁-C₆alkoxy; and

(ii) 2-indanyl, substituted with 0, 1 or 2 substituents independently selected from fluoro, chloro, hydroxy, methyl, ethyl, methoxy, ethoxy, mono-, di- and tri-fluoromethyl, and mono-, di- and tri-fluoromethoxy;

R⁴ is C₁-C₆alkyl, haloC₁-C₆alkyl, fluoro or chloro;

R⁷ is hydrogen or C₁-C₆alkyl; and

R⁹ represents from 0 to 5 substituents independently chosen from hydrogen, halogen, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆haloalkyl, and C₁-C₆haloalkoxy.

36. A compound according to Claim 34, wherein the compound is N-(2-Fluorobenzyl)-N-indan-2-yl-2-(4-o-tolyl-6,7-dihydro-4H-thieno[3,2-c]pyridin-5-yl)-acetamide.

37. A compound of or salt Claim 1, having a enantiomeric excess of at least 90%.

38. A pharmaceutical composition comprising at least one compound or salt according to Claim 1, or a prodrug or hydrate thereof, in combination with a physiologically acceptable carrier or excipient.

39. A compound or salt according to Claim 1, wherein the compound exhibits an IC₅₀ of 100 nM or less in a standard *in vitro* C5a receptor-mediated chemotaxis or calcium mobilization assay.

40. A compound or salt according to Claim 1, wherein the compound exhibits less than 5% agonist activity in a GTP binding assay.

41. A method for inhibiting signal-transducing activity of a cellular C5a receptor, comprising contacting a cell expressing a C5a receptor with at least one compound or salt according to Claim 1, and thereby reducing signal transduction by the C5a receptor.

42. A method according to Claim 41, wherein the cell is contacted *in vivo* in an animal.

43. A method according to Claim 42, wherein the animal is a human.

44. A method of inhibiting binding of C5a to C5a receptor *in vitro*, the method comprising contacting C5a receptor with at least one compound or salt according to Claim 1, under conditions and in an amount sufficient to detectably inhibit C5a binding to C5a receptor.

45. A method of inhibiting binding of C5a to C5a receptor in a human patient, comprising contacting cells expressing C5a receptor with at least one compound or salt according to Claim 1, in an amount sufficient to detectably inhibit C5a binding to cells expressing a cloned C5a receptor *in vitro*, and thereby inhibiting binding of C5a to the C5a receptor in the patient.

46. A method for treating a patient suffering from rheumatoid arthritis, psoriasis, cardiovascular disease, reperfusion injury, or bronchial asthma comprising administering to the patient a C5a receptor modulatory amount of a compound according to Claim 1.

47. A method for treating a patient suffering from stroke, myocardial infarction, atherosclerosis, ischemic heart disease, or ischemia-reperfusion injury comprising administering to the patient a C5a receptor modulatory amount of a compound according to Claim 1.

48. A method for inhibiting C5a receptor-mediated cellular chemotaxis, comprising contacting mammalian white blood cells with a C5a receptor modulatory amount of a compound or salt according to Claim 1.

49. A method for localizing C5a receptors in a tissue sample, comprising:
contacting the tissue sample containing C5a receptors with a detectably labeled compound according to Claim 1 under conditions that permit binding of the compound to C5a receptors; and
detecting the bound compound.

50. A packaged pharmaceutical preparation, comprising:
(a) a pharmaceutical composition according to Claim 38 in a container; and
(b) instructions for using the composition to treat a patient suffering from rheumatoid arthritis, psoriasis, cardiovascular disease, reperfusion injury, or bronchial asthma.

51. A packaged pharmaceutical preparation
(a) a pharmaceutical composition according to Claim 38 in a container; and
(b) instructions for using the composition to treat stroke, myocardial infarction, atherosclerosis, ischemic heart disease, or ischemia-reperfusion injury.

52. A pharmaceutical composition according to Claim 38, wherein the pharmaceutical composition is formulated as an injectible fluid, an aerosol, a cream, a gel, a pill, a capsule, a syrup, or a transdermal patch.